

# Patients' beliefs about medicine are associated with early thiopurine discontinuation in patients with inflammatory bowel diseases

Mark M.T.J. Broekman<sup>a</sup>, Marieke J.H. Coenen<sup>b</sup>, Geert J. Wanten<sup>a</sup>, Corine J. van Marrewijk<sup>b</sup>, Wietske Kievit<sup>c</sup>, Olaf H. Klungel<sup>d</sup>, André L.M. Verbeek<sup>c</sup>, Dennis R. Wong<sup>e</sup>, Piet M. Hooymans<sup>e</sup>, Henk-Jan Guchelaar<sup>f</sup>, Hans Scheffer<sup>b</sup>, Luc J.J. Derijks<sup>h</sup>, Marcel L. Bouvy<sup>d,g</sup> and Dirk J. de Jong<sup>a</sup>

**Background** Patients' beliefs about medicine may either reflect the necessity for treatment or concerns regarding the treatment. We explored the extent to which these beliefs have an effect on thiopurine metabolite levels and premature discontinuation in patients with inflammatory bowel disease (IBD).

**Patients and methods** Patients enrolled in the 'Thiopurine response Optimization by Pharmacogenetic testing in Inflammatory Bowel Disease Clinics' (TOPIC) trial were asked to complete the Beliefs about Medicine Questionnaire (BMQ) 4 weeks after thiopurine initiation. The BMQ measures perceptions about treatment necessity and concerns. On the basis of the necessity and concern scores, patients can be categorized as accepting, ambivalent, indifferent, or skeptical. The thiopurine discontinuation rates for these belief subgroups were compared by Kaplan-Meier curves. Furthermore, clinical response and metabolite levels were compared between the belief subgroups.

**Results** A total of 767 patients with IBD started thiopurine treatment, of whom 576 (75%) completed the BMQ. Patients could be classified as accepting (34%), indifferent (17%), ambivalent (34%), or skeptical (15%). Compared with patients in the accepting group (discontinuation rate 22%), patients with an indifferent (35%;  $P=0.02$ ), ambivalent (37%;  $P<0.01$ ), or skeptical belief (54%;  $P<0.01$ ) had higher thiopurine discontinuation rates. No differences were observed in the steady-state thiopurine metabolite levels between the different belief subgroups.

**Conclusion** Patients with a low perceived treatment necessity or high concerns toward IBD treatment were more likely to discontinue thiopurine treatment prematurely. Extra attention toward these patients might prevent premature discontinuation. Eur J Gastroenterol Hepatol 00:000–000

Copyright © 2017 The Author(s). Published by Wolters Kluwer Health, Inc.

## Introduction

Thiopurines are seen as the cornerstone in the maintenance treatment of inflammatory bowel disease (IBD). Although thiopurines are effective in the treatment of ulcerative

colitis (UC) and Crohn's disease (CD), up to 40% of patients discontinue treatment within 1 year [1,2]. Several factors may result in early treatment discontinuation, such as occurrence of side effects or lack of effectiveness. Apart from the actual occurrence of side effects or the lack of effectiveness, patients' beliefs about the necessity for the treatment (e.g. the personal need for the prescribed drug) and concerns regarding potential drawbacks of treatment (e.g. side effects, long-term effects, or dependence) may influence the motivation to start and continue medication [3,4].

Patients with IBD have relatively high levels of necessity, which suggests a high need for medical treatment [3,5]. Nevertheless, nonadherence to medication is reported in 30–60% of patients [3,6]. This results in an increased flare risk and a lower quality of life [6]. Nonadherence to thiopurines might seduce the attending physician to an unjustified switch to expensive biologic drugs in an effort to control disease, thereby placing an increased burden on healthcare resources. In IBD, as in other diseases, nonadherence is more frequently encountered in patients with stronger concerns or lower perceived necessity of treatment [7–10].

Evaluation of treatment adherence is a challenge. Objective methods to assess treatment adherence such as drug tests in blood or urine are cumbersome and not available for all drugs; therefore, adherence is frequently assessed by indirect

European Journal of Gastroenterology & Hepatology 2017, 00:000–000

**Keywords:** adherence, beliefs about medicine, inflammatory bowel disease, thiopurine

<sup>a</sup>Department of Gastroenterology, Radboud Institute for Molecular Life Sciences, Departments of <sup>b</sup>Human Genetics, <sup>c</sup>Health Evidence, Radboud Institute for Health Sciences, Radboud University Medical Center, Nijmegen, <sup>d</sup>Department of Pharmacoepidemiology and Pharmacotherapy, Utrecht University, Utrecht, <sup>e</sup>Department of Clinical Pharmacy, Pharmacology and Toxicology, Zuyderland Medical Center, Sittard-Geleen, <sup>f</sup>Department of Clinical Pharmacy and Toxicology, Leiden University Medical Center, <sup>g</sup>SIR institute for Pharmacy Practice and Policy, Leiden and <sup>h</sup>Department of Clinical Pharmacy, Máxima Medical Center, Veldhoven, The Netherlands

Correspondence to Mark M.T.J. Broekman, MD, Department of Gastroenterology, Radboud Institute for Molecular Life Sciences, Radboud University Medical Center, PO Box 9101, 6500 HB Nijmegen, The Netherlands

Tel: +31 24 3614760; fax: +31 24 3540103; e-mail: broekmanmark@gmail.com

**Received** 12 September 2017 **Accepted** 11 October 2017

Supplemental Digital Content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website, [www.eurojgh.com](http://www.eurojgh.com).

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

and subjective methods [11,12]. Unfortunately, significant discrepancies exist between self-reported adherence and objective assays [13]. In thiopurine treatment, assessment of the thiopurine metabolites 6-methylmercaptopurine ribonucleotides (6-MMPR) and 6-thioguanine nucleotide (6-TGN) levels is becoming increasingly available in daily clinic for optimization and evaluation of thiopurine treatment. The presence of low levels of 6-MMPR and 6-TGN may suggest noncompliance and can therefore be used as a tool to assess treatment adherence [14]. Manipulation of adherence by taking pills just before a hospital visit is less relevant with thiopurines because it takes several weeks to achieve steady-state metabolite levels.

Identification of patients at risk for nonadherence or early treatment discontinuation would be useful in designing a targeted approach to improve adherence. Apart from the influence of personal beliefs on self-reported treatment adherence and factors associated with adherence, little is known on the effect of beliefs about medicine on steady-state thiopurine metabolite levels and, more importantly, how these beliefs may foresee treatment discontinuation [5,12]. To this end, we explored whether beliefs about IBD drugs could be associated with metabolite levels or premature treatment discontinuation of thiopurines.

## Patients and methods

### Patients

The current study is a post-hoc analysis of the Thiopurine response Optimization by Pharmacogenetic testing in Inflammatory bowel disease Clinics (TOPIC) trial. In the TOPIC trial, thiopurine-naïve patients with IBD were randomized for standard-of-care thiopurine dosing versus a genotype-based dose to evaluate the efficacy of pretreatment thiopurine S-methyltransferase genotyping on the occurrence of myelosuppression [15]. Patients were followed for  $20 \pm 6$  weeks. Physicians were free to use either azathioprine or mercaptopurine. The main exclusion criteria were previous use of thiopurines, laboratory test abnormalities, known thiopurine S-methyltransferase enzyme activity, or genotype. The study protocol of the TOPIC trial was approved by the Institutional review board of the Radboud University Medical Center. All patients provided written informed consent. The TOPIC trial was registered at clinicaltrials.gov, NCT00521950. The TOPIC trial complies with the Declaration of Helsinki. For a detailed description of the study design, patient selection, and patient data, we refer to Coenen *et al.* [15].

### Beliefs about Medicine Questionnaire

Patients participating in the TOPIC trial were asked to complete the Beliefs about Medicine Questionnaire (BMQ) and the Medication Adherence Report Scale (MARS) 4 weeks after thiopurine initiation. Personal needs and concerns regarding a treatment can be quantified with the BMQ [16]. The BMQ is validated in patients with various chronic diseases such as asthma, rheumatoid arthritis, diabetic, renal dialysis, chronic heart disease, IBD, and psychiatric diseases [16,17].

For this study, the BMQ included statements about IBD treatment in a random order in terms of the necessity (five questions; e.g. My future health depends on the use of my

IBD medication) and concerns (six questions; e.g. The use of these IBD drugs disrupt my life). Statements are scored on a Likert-type scale (ranging from 1 = strongly disagree to 5 = strongly agree). The sum of answers for the five statements about necessity leads to a score between 5 and 25. For concern statements, the sum of answers was divided by 6 (the number of statements) and multiplied by 5 to generate a score between 5 and 25. Plotting the necessity score at the y-axis and the concerns score at the x-axis, with a split at the midpoint, enables a classification into four belief subgroups (Table 1) [7].

### Self-reported adherence

Self-reported adherence was assessed using the five-item MARS questionnaire [16]. Questions are related to several statements concerning adherence (e.g. I forget to take my medication or I take fewer pills than prescribed). Answers were provided on a five-point Likert scale. The sum of answers was used as a subjective marker of adherence, with a maximum score of 25, which can be considered full (self-reported) adherence.

### 6-Methylmercaptopurine ribonucleotides and 6-thioguanine nucleotide metabolite measurements

By protocol, 6-TGN and 6-MMPR levels at week 8 were assessed at a routine check-up in the first 301 patients included in the TOPIC trial. 6-TGN and 6-MMPR levels were determined in red blood cells (RBCs) by high-performance liquid chromatography according to the Lennard method [15]. Thiopurine metabolite concentrations are presented in  $\text{pmol}/8 \times 10^8$  RBCs. For analyses with metabolite levels, we used data from patients without dose adjustments in the first 8 weeks. Subtherapeutic levels were defined as 6-TGN levels below  $230 \text{ pmol}/8 \times 10^8$  RBCs, whereas therapeutic levels were defined as 6-TGN levels between 230 and  $490 \text{ pmol}/8 \times 10^8$  RBCs [18].

### Clinical response

Clinical response was evaluated using the Harvey–Bradshaw Index in patients with CD and the partial Mayo score in patients with UC. Disease activity was measured at baseline, week 0, and at the end of the clinical trial, week 20 [19]. Treatment response was defined as a reduction of three or more points at week 20 compared with week 0 on either the Harvey–Bradshaw Index or the partial Mayo score.

### Data analysis

For this study, all patients who completed the BMQ were included. One missing answer per subscale (necessity, concerns, or self-reported adherence part) was accepted; in that case, the

**Table 1.** Classification of patients into four belief subgroups on the basis of the Beliefs about Medicine Questionnaire score on the necessity and concerns regarding inflammatory bowel disease treatment

Belief subgroup	Necessity score	Concern score
Accepting	High	Low
Ambivalent	High	High
Indifferent	Low	Low
Skeptical	Low	High

A high score on both the necessity and the concern subscale is defined as a score of more than 15 points and a low score is defined as 15 points or lower.

missing value was replaced by the mean of the other answers on that subscale. Patients were excluded from this study if more than one answer per subscale was missing. Scores are presented as mean  $\pm$  SD, unless mentioned otherwise. Differences between respondents and nonrespondents were compared using a parametric and a nonparametric test, where appropriate. Necessity and concerns scores were compared for CD and UC using an independent-samples *t*-test. In the same way, necessity and concerns scores were compared between patients who continued and discontinued their thiopurine during the 5 months after treatment initiation. Treatment discontinuation was defined as a patient who stopped the initial prescribed thiopurine within 5 months. Dose reductions and temporary interruptions were accepted, provided that the interruption was not longer than 1 month [20]. Kaplan–Meier curves were plotted to illustrate the time to discontinuation for the four belief subgroups and the log-rank test was used to compare discontinuation rates. Self-reported adherence rates were compared between the different belief subgroups using the Kruskal–Wallis test. The Kruskal–Wallis test was used to compare 6-MMPR and 6-TGN levels between the different belief subgroups and the  $\chi^2$ -test was used to compare the number of patients with 6-TGN levels within the therapeutic range. The degree of correlation between self-reported adherence scores and metabolite levels was examined by Spearman's correlation coefficient. Differences in treatment response for the belief subgroups were compared using the chi-squared test. A *P*-value of less than 0.05 was considered significant. All analyses were carried out in SPSS, version 22.0.0.1 (SPSS Inc., Chicago, Illinois, USA).

## Results

### Patients

Of the 796 randomized patients in the TOPIC trial, 767 patients actually started with thiopurines and 601 (78%) returned the BMQ. A total of 25 returned questionnaires were excluded because of more than one missing answer in one of the three subscales, leaving 576 (75%) patients, the respondents who successfully completed the questionnaires for further analysis. Patients who did not return the questionnaire (*n* = 166) or who were excluded because of missing answers (*n* = 25) were classified as nonrespondents. The characteristics of respondents and nonrespondents are shown in Table 2.

### Beliefs about medicine

Patients with IBD scored ( $17.1 \pm 4.2$ ) on the necessity subscale. Necessity scores were higher in patients with UC ( $17.8 \pm 4.2$ ) than CD ( $16.6 \pm 4.1$ ) (*P* < 0.01). Concern scores were not different between UC ( $15.6 \pm 3.4$ ) and CD ( $15.1 \pm 3.4$ ) (*P* = 0.07). Categorization of patients according to the combination of necessity and concerns score showed that 34% of the patients could be defined as accepting, 34% as ambivalent, 17% as indifferent, and 15% as skeptical (Fig. 1).

### Thiopurine discontinuation rates

Of the 576 respondents, 197 (34%) discontinued the initial prescribed thiopurine within the first 5 months. Patients who discontinued treatment had higher concerns ( $16.3 \pm 3.6$  vs.  $14.8 \pm 3.2$ , *P* < 0.01) and lower necessity scores ( $16.1 \pm 4.9$

vs.  $17.7 \pm 3.6$ , *P* < 0.01) than patients who continued thiopurine treatment. Compared with patients in the accepting group (discontinuation rate 22%), patients classified as either indifferent (discontinuation rate 35%; *P* = 0.02), ambivalent (discontinuation rate 37%; *P* < 0.01), or skeptical (discontinuation rate 54%; *P* < 0.01) all had higher discontinuation rates (Fig. 2). Moreover, discontinuation rates of patients with a skeptical profile were also higher compared with patients classified as either indifferent or ambivalent (*P* < 0.01).

### Self-reported adherence

Self-reported adherence scores, measured with the MARS questionnaire, were very high, with a median score (inter-quartile range) of 25.0 (24.0–25.0). Self-reported adherence scores were not different between the belief subgroups (*P* = 0.88). 6-MMPR metabolite levels (*P* = 0.29) and 6-TGN metabolite levels (*P* = 0.34) were not correlated with the self-reported adherence.

### Thiopurine metabolite levels

Overall, 196 (65%) of the 301 patients in whom the week eight 6-MMPR and 6-TGN levels were assessed were considered as steady state (i.e. had no dose adjustments during the first 8 weeks) and were included for further analyses [21]. The number of patients per belief subgroup with 6-TGN levels within the therapeutic range varied between 43 and 48%. 6-MMPR (*P* = 0.99), 6-TGN (*P* = 0.97) levels and the percentage of patients within the therapeutic range (*P* = 0.97) were not different between the four belief subgroups (Table 3).

### Clinical response

Clinical response was known for 338 (58.7%) of the 576 respondents. Categorization of these 388 patients in belief subgroups showed that 35% of the patients could be classified as accepting, 35% as ambivalent, 16% as indifferent, and 14% as skeptical, which is in line with the numbers of the entire group (Fig. 1). Treatment response rates per belief subgroup were 26% (30 out of 117) for accepting patients, 26% (31 out of 121) in ambivalent patients, 17% (nine out of 54) in indifferent patients, and 34% (16 of 47) in skeptical patients (Fig. 3). Treatment response was not different between the four belief subgroups (*P* = 0.26).

## Discussion

The present study shows that patients with IBD who perceive a low necessity or high concerns for IBD treatment were more likely to discontinue thiopurine treatment prematurely. Categorization into four belief subgroups showed that most patients could be classified as accepting or ambivalent. We found no association between belief subgroups and 6-MMPR and 6-TGN metabolite levels.

The proportion of patients per belief subgroup corroborates with those found in previous studies in IBD patients [3,22]. The relatively high scores of patients toward the necessity for treatment probably follow from a high level of disease burden in patients with IBD [23,24]. Application of the BMQ in diseases with a lower disease

**Table 2.** Characteristics of respondents and nonrespondents


	Total (n = 767) [n (%)]	Respondents (n = 576) [n (%)]	Nonrespondents (N = 191) [n (%)]	P-value
Age [median (IQR)] (years)	41 (27–53)	43 (28–55)	37 (24–47)	<0.01
Sex: female	421 (55)	323 (56)	98 (51)	0.25
Disease type, CD <sup>a</sup>	463 (60)	336 (59)	127 (66)	0.05
Disease duration [mean (SD)] (years)	5.3 (8.2)	5.7 (8.6)	4.1 (6.9)	0.01
Baseline disease severity <sup>b</sup>				
CD (HBI) [mean (SD)]	3.5 (3.0)	3.2 (2.7)	4.3 (3.8)	0.01
UC (partial Mayo score) [mean (SD)]	3.8 (1.7)	3.8 (1.7)	4.3 (1.7)	0.10
Type of thiopurine, AZA	494 (64)	380 (66)	114 (60)	0.12
Discontinuation before week 20	296 (39)	197 (34)	99 (52)	<0.01

The nonrespondents group is composed of patients who did not return the questionnaire (n = 166) and patients with more than 1 missing answer in one of the three subscales (n = 25).


AZA; azathioprine, CD; Crohn's disease, HBI; Harvey–Bradshaw Index, IQR, interquartile range; UC; ulcerative colitis.

<sup>a</sup>Seven patients were diagnosed with indeterminate colitis.

<sup>b</sup>HBI was known for 349 patients with CD and the partial mayo score was known for 251 UC patients.



**Fig. 1.** Categorization of patients in four belief subgroups on the basis of their necessity and concerns score, including the discontinuation rate per belief subgroup.



**Fig. 2.** Kaplan–Meier curve in which treatment discontinuation is plotted for the four belief subgroups: accepting group (n = 198), ambivalent group (n = 198), indifferent group (n = 96), and skeptical group (n = 84).

burden, such as atrial fibrillation, indeed results in lower necessity scores than thrombosis, which has more impact [25]. The higher necessity scores of UC patients compared with patients with CD might be explained by the fact that UC exacerbations with frequent bloody diarrhea may be perceived as more severe compared with CD, which can be more subtle in presentation [26]. The fact that a third of the patients with IBD scored low on necessity for treatment may have resulted from the cyclical disease manifestation

with episodes of no or only minimal disease symptoms [12].

A novel aspect of this study is that we correlated BMQ scores with treatment discontinuation rates. To date, studies using the BMQ in patients with IBD mainly focused on the exploration of factors associated with drug adherence [12,27]. Knowledge of who discontinues treatment is of high relevance, given the limited alternative options in the treatment of IBD. We showed that IBD patients with high concerns or low needs are at risk for premature discontinuation of thiopurines in the first months of treatment. Moreover, the combination of both even further increases this risk. Our findings corroborate studies in patients with IBD that showed a positive relation between the patients' necessity and medication adherence and a negative relation in case of high concerns [12]. In the current study, the BMQ was collected 4 weeks after thiopurine initiation, which is in line with comparable study designs [25,28]. Although studies showed that patients' beliefs are considered stable over time, the occurrence of side effects within the first weeks may have resulted in increased concern feelings at the time that the questionnaire was completed [29]. Identification of patients at risk for treatment discontinuation in an early stage might help in the design of strategies to prevent them from discontinuing treatment. Special attention should be paid to patients with high concerns as within the spectrum of IBD drugs, patients consider thiopurines as drugs with the highest risks as a result of fear of side effects and long-term drug-induced comorbidities [30]. This leads to significantly lower adherence rates for thiopurine use [31]. A comprehensive education at start and follow-up phone calls by an IBD nurse or pharmacist seem to be a promising approach to evaluate and anticipate the negative perceptions and expectations toward thiopurine treatment [6,32–34].


This study is unique as thiopurine metabolite levels were used as an indicator for treatment adherence in conjunction with self-reported adherence scores. Most studies only used self-reported adherence or pill count as a proxy for treatment adherence [3,9]. In our study, patients scored very high on self-reported adherence, whereas the number of patients in the therapeutic range varied around 45%. Previous studies reported lower self-reported adherence scores, suggesting that self-reported adherence scores in this study might be biased because of the setting of a clinical trial [3,7,9,24]. Only a few studies have compared self-reported adherence with thiopurine metabolite levels measured in blood. In one



**Table 3.** Steady-state week eight 6-methylmercaptopurine ribonucleotides, 6-thioguanine nucleotide levels, and percentage of patients within the therapeutic range per belief subgroup

Belief subgroup	6-MMPR (pmol/8 × 10 <sup>8</sup> RBCs)	6-TGN (pmol/8 × 10 <sup>8</sup> RBCs)	Patients within the therapeutic range (%)
Accepting ( <i>n</i> = 76; 38%)	3068 (918–7352)	259 (204–381)	47
Ambivalent ( <i>n</i> = 63; 32%)	3141 (681–7363)	271 (196–343)	48
Indifferent ( <i>n</i> = 36; 18%)	3639 (867–6219)	271 (195–378)	44
Skeptical ( <i>n</i> = 21; 11%)	4101 (613–8024)	264 (171–508)	43

Week 8 metabolite levels were presented as median (interquartile range). Therapeutic range was defined as 6-TGN levels between 230 and 490 pmol/8 × 10<sup>8</sup> RBCs. 6-MMPR (*P* = 0.99), 6-TGN levels (*P* = 0.97), and therapeutic range (*P* = 0.97) were not different between the different belief subgroups. 6-MMPR, 6-methylmercaptopurine ribonucleotides; RBC, red blood cell; 6-TGN, 6-thioguanine nucleotide.

**Fig. 3.** Percentage of patients with a reduction of three or more points on the Harvey-Bradshaw Index or partial Mayo score after 20 weeks for the four belief subgroups. Treatment response was not different between subgroups (*P* = 0.26).

of these studies, self-reported nonadherence was limited to 10% of thiopurine users, whereas in practice, only 14% had 6-TGN levels within the therapeutic range [13]. It must be stressed that many genetic and environmental factors influence thiopurine metabolism, which limits analyses between necessity or concern scores and metabolite levels [35]. This underscores the relevance, but also the difficulty, of more objective outcome measures. We found no association between the four belief subgroups and 6-MMPR and 6-TGN metabolite levels at week 8. Furthermore, there were no differences in treatment response between the belief subgroups, which seems logical, given the similar 6-MMPR and 6-TGN levels across the belief subgroups. This indicates that, at least in the short term, the BMQ does not allow to predict adherence or treatment response in patients with IBD, but only who will discontinue treatment.

A limitation of our study is that obtaining metabolite levels at week 8 might be too early in the follow-up to identify differences in levels between the belief subgroups. Nonadherence risk increases with length of treatment, whereas in this study, patients just started treatment and might still be motivated to take their pills [36]. This is supported by a study in 65 patients with IBD with an average thiopurine usage of almost 3 years, in which metabolite levels were significantly lower in patients with a low self-reported adherence [37]. Furthermore, we only used data from patients with steady-state week 8 metabolite levels. Patients with dose adjustments, frequently because of side

effects, were excluded. Importantly, proportions of the four belief subgroups of patients with steady-state metabolite levels merely corresponded with numbers of the entire study group, which makes selection bias unlikely. Another limitation is that selection bias occurred, given the differences between respondents and nonrespondents in age, disease duration, and discontinuation rates. The latter can partially be explained by patients who discontinued treatment before week 4 and subsequently did not return the questionnaire [38,39]. Finally, questions in the BMQ and MARS questionnaire contained the term 'IBD treatment' rather than 'thiopurine treatment' to avoid confusion with complicated chemical drug names (azathioprine or mercaptopurine) or brand names. Therefore, it might be that patients did not answer the questions with reference to thiopurine use. However, we foresee that this influence is minimal as patients were asked to participate in a trial that focused on the initiation of thiopurines.

## Conclusion

The use of the BMQ helps to identify IBD patients who are likely to discontinue thiopurines in the first months of treatment. Attention and education in the consulting room or by telephone should be given to patients with low needs and high concerns toward IBD treatment to prevent premature discontinuation.

## Acknowledgements

The authors are deeply indebted to the participants of the TOPIC trial. Members of the TOPIC recruitment team responsible for patient recruitment and collection of clinical data are listed in Supplementary File 1, Supplemental Digital Content 1, <http://links.lww.com/EJGH/A237>. Compensation was given to the members of the recruitment team for additional biochemical measurements and examinations that had to be performed for the TOPIC study. Furthermore, we thank Mariëlle Maas, Miet Fiddelaers, Milevis Reitsma, Leonie Peters, and Jean Cilissen from the department of Clinical Pharmacy and Toxicology, Zuyderland Medical Center, Sittard-Geleen, The Netherlands, for technical assistance with metabolite measurement and Debbie Heinen, Marjolein M.J. van Donkelaar, Freshteh Golestani, Marlies E. de Vos, J.G. Angelen M. Heister, Doménique M.W. Nijsten, Mascha M.V.A.P. Schijvenaars, and Martine E.C. Cranen from the department of Human Genetics, Radboud university medical center, Nijmegen, The Netherlands, for their support in data management. We thank dr. Sita H. Vermeulen, department for Health Evidence, Radboud Institute for Health Sciences, Radboud University Medical

Center, Nijmegen, The Netherlands, and professor Barbara Franke, department of Human Genetics, Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center, Nijmegen, The Netherlands, for their contribution to the design of the TOPIC trial. Finally, we thank professor Joost P.H. Drenth from the Department of Gastroenterology, Radboud University Medical Center, Nijmegen, The Netherlands, for intellectual contribution to the content of the manuscript.

M.B., O.K., M.C., M.L.B., D.J. contributed to the study design; M.B., C.M., M.C., D.W., P.H., D.J., G.W. carried out data collection; M.B., W.K., M.L.B. carried out data analysis; M.B., W.K., D.J., G.W., M.L.B. carried out interpretation of data; M.B., G.W., D.J., M.C., M.L.B. contributed in writing of the manuscript; H.G., A.V., H.S., L.D., O.K., W.K. carried out critical revision of the manuscript for important intellectual content; and D.J., M.C. supervised the study.

The TOPIC trial was funded by The Netherlands Organisation for Health Research and Development (grant number 94507606) and the participating institutes.

### Conflicts of interest

There are no conflicts of interest.

### References

- 1 Terdiman JP, Gruss CB, Heidelbaugh JJ, Sultan S, Falck-Ytter YT. AGA Institute Clinical Practice and Quality Management Committee. American Gastroenterological Association Institute guideline on the use of thiopurines, methotrexate, and anti-TNF-alpha biologic drugs for the induction and maintenance of remission in inflammatory Crohn's disease. *Gastroenterology* 2013; 145:1459–1463.
- 2 Jharap B, Seinen ML, de Boer NK, van Ginkel JR, Linskens RK, Kneppelhout JC, et al. Thiopurine therapy in inflammatory bowel disease patients: analyses of two 8-year intercept cohorts. *Inflamm Bowel Dis* 2010; 16:1541–1549.
- 3 Horne R, Parham R, Driscoll R, Robinson A. Patients' attitudes to medicines and adherence to maintenance treatment in inflammatory bowel disease. *Inflamm Bowel Dis* 2009; 15:837–844.
- 4 Vangeli E, Bakhshi S, Baker A, Fisher A, Bucknor D, Mrowietz U, et al. A Systematic review of factors associated with non-adherence to treatment for immune-mediated inflammatory diseases. *Adv Ther* 2015; 32:983–1028.
- 5 Kim SB, Kim KO, Jang BI, Kim ES, Cho KB, Park KS, et al. Patients' beliefs and attitudes about their treatment for inflammatory bowel disease in Korea. *J Gastroenterol Hepatol* 2016; 31:575–580.
- 6 Greenley RN, Kunz JH, Walter J, Hommel KA. Practical strategies for enhancing adherence to treatment regimen in inflammatory bowel disease. *Inflamm Bowel Dis* 2013; 19:1534–1545.
- 7 Menckeborg TT, Bouvy ML, Bracke M, Kaptein AA, Leufkens HG, Raaijmakers JA, et al. Beliefs about medicines predict refill adherence to inhaled corticosteroids. *J Psychosom Res* 2008; 64:47–54.
- 8 Chapman SC, Barnes N, Barnes M, Wilkinson A, Hartley J, Piddock C, et al. Changing adherence-related beliefs about ICS maintenance treatment for asthma: feasibility study of an intervention delivered by asthma nurse specialists. *BMJ Open* 2015; 5:e007354.
- 9 Ediger JP, Walker JR, Graff L, Lix L, Clara I, Rawsthorne P, et al. Predictors of medication adherence in inflammatory bowel disease. *Am J Gastroenterol* 2007; 102:1417–1426.
- 10 Kamperidis N, Goodhand JR, Chowdhury FA, Koodun Y, Direzke NC, Naik S, et al. Factors associated with nonadherence to thiopurines in adolescent and adult patients with inflammatory bowel disease. *J Pediatr Gastroenterol Nutr* 2012; 54:685–689.
- 11 Lakatos PL. Prevalence, predictors, and clinical consequences of medical adherence in IBD: how to improve it? *World J Gastroenterol* 2009; 15:4234–4239.
- 12 Horne R, Chapman SC, Parham R, Freemantle N, Forbes A, Cooper V. Understanding patients' adherence-related beliefs about medicines prescribed for long-term conditions: a meta-analytic review of the Necessity-Concerns Framework. *PLoS one* 2013; 8:e80633.
- 13 Hommel KA, Davis CM, Baldassano RN. Objective versus subjective assessment of oral medication adherence in pediatric inflammatory bowel disease. *Inflamm Bowel Dis* 2009; 15:589–593.
- 14 Coskun M, Steenholdt C, de Boer NK, Nielsen OH. Pharmacology and optimization of thiopurines and methotrexate in inflammatory bowel disease. *Clin Pharmacokinet* 2016; 55:257–274.
- 15 Coenen MJ, de Jong DJ, van Marrewijk CJ, Derijks LJ, Vermeulen SH, Wong DR, et al. Identification of patients with variants in TPMT and dose reduction reduces hematologic events during thiopurine treatment of inflammatory bowel disease. *Gastroenterology* 2015; 149:907–917e7.
- 16 Horne R, Weinman J. Patients' beliefs about prescribed medicines and their role in adherence to treatment in chronic physical illness. *J Psychosom Res* 1999; 47:555–567.
- 17 Horne R, Weinman J, Harkins M. The beliefs about medicines questionnaire: The development and evaluation of a new method for assessing the cognitive representation of medication. *Psychol Health* 1999; 14:1–24.
- 18 Gilissen LP, Wong DR, Engels LG, Bierau J, Bakker JA, Paulussen AD, et al. Therapeutic drug monitoring of thiopurine metabolites in adult thiopurine tolerant IBD patients on maintenance therapy. *J Crohns Colitis* 2012; 6:698–707.
- 19 Lewis JD, Chuai S, Nessel L, Lichtenstein GR, Abera FN, Ellenberg JH. Use of the noninvasive components of the Mayo score to assess clinical response in ulcerative colitis. *Inflamm Bowel Dis* 2008; 14:1660–1666.
- 20 Broekman M, Coenen MJH, van Marrewijk CJ, Wanten GJA, Wong DR, Verbeek ALM, et al. More dose-dependent side effects with mercaptopurine over azathioprine in ibd treatment due to relatively higher dosing. *Inflamm Bowel Dis* 2017; 23:1873–1881.
- 21 Broekman MM, Wong DR, Wanten GJ, Roelofs HM, van Marrewijk CJ, Klungel OH, et al. The glutathione transferase Mu null genotype leads to lower 6-MMPR levels in patients treated with azathioprine but not with mercaptopurine. *Pharmacogenomics J* 2017. [Epub ahead of print].
- 22 Moshkovska T, Stone MA, Clatworthy J, Smith RM, Bankart J, Baker R, et al. An investigation of medication adherence to 5-aminosalicylic acid therapy in patients with ulcerative colitis, using self-report and urinary drug excretion measurements. *Aliment Pharmacol Ther* 2009; 30 (11–12):1118–1127.
- 23 Cosnes J, Gower-Rousseau C, Seksik P, Cortot A. Epidemiology and natural history of inflammatory bowel diseases. *Gastroenterology* 2011; 140:1785–1794.
- 24 Sjolander M, Eriksson M, Glader EL. The association between patients' beliefs about medicines and adherence to drug treatment after stroke: a cross-sectional questionnaire survey. *BMJ Open* 2013; 3:e003551.
- 25 Verhoef TI, Redekop WK, Bouvy ML, Dorenbos B, Karwar Z, van Schie RM, et al. Beliefs about medicines in Dutch acenocoumarol and phenprocoumon users. *Br J Clin Pharmacol* 2014; 78:422–429.
- 26 Targownik LE, Sexton KA, Bernstein MT, Beatie B, Sargent M, Walker JR, et al. The Relationship among perceived stress, symptoms, and inflammation in persons with inflammatory bowel disease. *Am J Gastroenterol* 2015; 110:1001–1012. quiz 1013.
- 27 Goodhand JR, Kamperidis N, Sirwan B, Macken L, Tshuma N, Koodun Y, et al. Factors associated with thiopurine non-adherence in patients with inflammatory bowel disease. *Aliment Pharmacol Ther* 2013; 38:1097–1108.
- 28 Linn AJ, van Weert JC, Schouten BC, Smit EG, van Bodegraven AA, van Dijk L. Words that make pills easier to swallow: a communication typology to address practical and perceptual barriers to medication intake behavior. *Patient Prefer Adherence* 2012; 6:871–885.
- 29 Unni E, Shiyabola OO, Farris KB. Change in medication adherence and beliefs in medicines over time in older adults. *Glob J Health Sci* 2015; 8:51207.
- 30 Vaucher C, Maillard MH, Froehlich F, Burnand B, Michetti P, Pittet V. Patients and gastroenterologists' perceptions of treatments for inflammatory bowel diseases: do their perspectives match? *Scand J Gastroenterol* 2016; 51:1056–1061.
- 31 Campos S, Portela F, Sousa P, Sofia C. Inflammatory bowel disease: adherence to immunomodulators in a biological therapy era. *Eur J Gastroenterol Hepatol* 2016; 28:1313–1319.
- 32 Clifford S, Barber N, Elliott R, Hartley E, Horne R. Patient-centred advice is effective in improving adherence to medicines. *Pharm World Sci* 2006; 28:165–170.
- 33 Bouvy ML, Heerdink ER, Urquhart J, Grobbee DE, Hoes AW, Leufkens HG. Effect of a pharmacist-led intervention on diuretic compliance in heart failure patients: a randomized controlled study. *J Card Fail* 2003; 9:404–411.

- 34 Brown MT, Bussell JK. Medication adherence: WHO cares? *Mayo Clin Proc* 2011; 86:304–314.
- 35 Derijks LJ, Wong DR. Pharmacogenetics of thiopurines in inflammatory bowel disease. *Curr Pharm Des* 2010; 16:145–154.
- 36 Mantzaris GJ, Roussos A, Kalantzis C, Koilakou S, Raptis N, Kalantzis N. How adherent to treatment with azathioprine are patients with Crohn's disease in long-term remission? *Inflamm Bowel Dis* 2007; 13:446–450.
- 37 Bokemeyer B, Teml A, Roggel C, Hartmann P, Fischer C, Schaeffeler E, *et al.* Adherence to thiopurine treatment in out-patients with Crohn's disease. *Aliment Pharmacol Ther* 2007; 26:217–225.
- 38 Lamberts MP, Den Ouden BL, Gerritsen JJ, Roukema JA, Westert GP, Drenth JP, *et al.* Prospective multicentre cohort study of patient-reported outcomes after cholecystectomy for uncomplicated symptomatic cholecystolithiasis. *Br J Surg* 2015; 102:1402–1409.
- 39 Multone E, Vader JP, Mottet C, Schoepfer A, Fournier N, Burnand B, *et al.* Characteristics of non-responders to self-reported questionnaires in a large inflammatory bowel disease cohort study. *Scand J Gastroenterol* 2015; 50:1348–1356.